



This is the print version of the [Skeptical Science](http://sks.to/co2up) article '[CO2 is not increasing](http://sks.to/co2up)', which can be found at <http://sks.to/co2up>.

Are CO2 levels increasing?

What The Science Says:

Currently, humans are emitting around 29 billion tonnes of carbon dioxide into the atmosphere per year. Around 43% remains in the atmosphere - this is called the 'airborne fraction'. The rest is absorbed by vegetation and the oceans. While there are questions over how much the airborne fraction is increasing, it is clear that the total amount of CO₂ in the atmosphere is increasing dramatically. Current CO₂ levels are the highest in 15 million years.

Climate Myth: CO2 is not increasing

"...there is the contention by Wolfgang Knorr of the Department of Earth Sciences at the University of Bristol in England that carbon dioxide levels in the atmosphere are about where they were 160 years ago." ([as quoted by Ken Ward Jr.](#))

The 'airborne fraction' refers to the amount of human CO₂ emissions remaining in the atmosphere. Approximately 43% of our CO₂ emissions stay in the atmosphere with the rest being absorbed by carbon sinks. But is the airborne fraction increasing? A paper published in November 2009 found no statistically significant trend ([Knorr 2009](#)). Anthony Watts labeled this result the "[Bombshell from Bristol](#)" - A potentially devastating result for anthropogenic global warming. Was it such a shock? The 2007 IPCC verdict on the airborne fraction was "*There is yet no statistically significant trend in the CO₂ growth rate since 1958 This 'airborne fraction' has shown little variation over this period.*" ([IPCC AR4](#)) I'm not sure the move from "not much happening" "to "still not much happening" warrants the label "bombshell".

The airborne fraction is calculated from the rate of human CO₂ emissions and changes in atmospheric CO₂ concentration. The global increase in atmospheric CO₂ has been directly measured since 1959 and can be calculated from ice cores for earlier periods. Primarily, CO₂ emissions come from fossil fuel combustion with a lesser contribution from land use changes. Fossil fuel combustion is calculated from international energy statistics. CO₂ emissions from land-use changes are more difficult to estimate and come with greater uncertainty. Land use emissions are estimated using deforestation and other land-use data, fire observations from space and carbon cycle modeling.

There have been several recent studies determining the airborne fraction. [Trends in the sources and sinks of carbon dioxide \(Le Quere 2009\)](#) examines the airborne fraction from 1959 to 2008. This period was chosen as we have directly measured atmospheric CO₂ levels over this time. Fossil fuel emissions rose steadily in recent decades, contributing 8.7 ± 0.5 gigatonnes of carbon in 2008. This is 41% greater than fossil fuel emissions in 1990. CO₂ emissions from land use was estimated at 1.2 ± 0.4 gigatonnes of carbon in 2008. Note the proportionally higher uncertainty compared to fossil fuel emissions.

Over this period, an average of 43% of each year's CO₂ emissions remained in the atmosphere although there is much year-to-year variability. The noise in the airborne fraction was reduced by removing the variability associated with El Nino Southern Oscillation (ENSO) and volcanic activity. They found the airborne fraction increased by $3 \pm 2\%$ per decade. This is a slightly increasing trend although only barely statistically significant .

[Knorr 2009](#) extends this analysis back to 1850 by combining direct CO₂ measurements from Mauna Loa and the South Pole with CO₂ data derived from Antarctic ice cores. This enabled Knorr to compare CO₂ emissions to atmospheric CO₂ levels for the past 150 years.

CO2 emissions versus CO2 levels




Figure 1: Observed increase atmospheric CO₂ derived from direct measurements, taking the average of Mauna Loa (Hawaii) and the South Pole (thin solid line) and two ice cores: Law Dome (dashed thin line) and Siple (thin dotted line). This is compared to total anthropogenic emissions (thick solid line) and 46% of total emissions (thick dashed line). ([Knorr 2009](#))

Knorr finds that since 1850, the airborne fraction has remained relatively constant. When CO₂ emissions were low, the amount of CO₂ absorbed by natural carbon sinks was correspondingly low. As human CO₂ emissions sharply increased in the 20th Century, the amount absorbed by nature increased correspondingly. The airborne fraction remained level at around 43%. The trend since 1850 is found to be $0.7 \pm 1.4\%$ per decade.

There are several differences in methodology between [Knorr 2009](#) and [Le Quere 2009](#). Knorr's result does not include the filtering for ENSO and volcanic activity employed by Le Quéré. However, when Knorr does include this filtering in his analysis, he finds a trend of $1.2 \pm 0.9\%$ per decade. This is smaller than Le Quere's result but is statistically significant.

Knorr also finds the 150 year trend while Le Quéré looks at the last 50 years. This may be significant. If the airborne fraction is increasing, it is possibly a recent phenomenon due to natural carbon sinks losing their absorption ability after becoming saturated. Several studies have found recent drops in the uptake of CO₂ by oceans ([Le Quere 2007](#), [Schuster 2007](#), [Park 2008](#)). However, with such a noisy signal, this is one question that will require more data before being more fully resolved.

Lastly, some perspective. There are still areas of uncertainty associated with the carbon cycle. Because of this uncertainty, scientists are currently debating whether the airborne fraction is steady at 43% or slightly increasing from 43%. Unfortunately, some skeptics use this uncertainty to hold the position that [the airborne fraction is closer to 0%](#).

Intermediate rebuttal written by John Cook

Update July 2015:

Here is a related lecture-video from [Denial101x - Making Sense of Climate Science Denial](#)

Additional videos from the MOOC

Andy Skuce: [The CO2 rise is man man-made](#)

Interviews with [various experts](#)

Expert interview with [Corinne Le Quéré](#)



Skeptical Science explains the science of global warming and examines climate misinformation through the lens of peer-reviewed research. The website won the Australian Museum 2011 Eureka Prize for the Advancement of Climate Change Knowledge. Members of the Skeptical Science team have authored peer-reviewed papers, a [college textbook on climate change](#) and the book [Climate Change Denial: Heads in the Sand](#). Skeptical Science content has been used in university courses, textbooks, government reports on climate change, television documentaries and numerous books.



The [Skeptical Science](#) website by Skeptical Science is licensed under a Creative Commons [Attribution 3.0 Unported License](#).