



This is the print version of the [Skeptical Science](#) article '[CO2 only causes 35% of global warming](#)', which can be found at <http://sks.to/35percent>.

How CO2 warming is driving climate

What The Science Says:

The Nature commentary by Penner et al. on which this argument is based actually says that on top of the global warming caused by carbon dioxide, other short-lived pollutants (such as methane and black carbon) cause an additional warming approximately 65% as much as CO₂, and other short-lived pollutants (such as aerosols) also cause some cooling. However, claiming that CO₂ has only caused 35% of global warming is a gross misinterpretation and misunderstanding of the paper.

Climate Myth: CO₂ only causes 35% of global warming

CO₂ does not account for even a majority of the warming seen over the past century. If other species accounted for 65% of historical warming that leaves only 35% for carbon dioxide. ([Doug Hoffman](#))

In August 2010, *Nature* published a [commentary by Penner et al.](#) which mainly focused on the uncertainty regarding the effect short-lived pollutants (such as aerosols and black carbon) have on the climate. As is often the case, many in the blogosphere misinterpreted and misunderstood the statements and conclusions in the commentary. Not surprisingly, the biggest misinterpretation related to the contribution of anthropogenic greenhouse gases to global warming. Below is the most misunderstood quote, with emphasis on the key word.

"Of the short-lived species, methane, tropospheric ozone and black carbon are key contributors to global warming, **augmenting** the radiative forcing of carbon dioxide by 65%. Others — such as sulphate, nitrate and organic aerosols — cause a negative radiative forcing, offsetting a fraction of the warming owing to carbon dioxide."

Numerous blogs have (mis)interpreted this statement to mean that carbon dioxide is only causing 35% as much global warming as previously believed. A more accurate reading of the quote is that certain short-lived pollutants cause warming **in addition to** carbon dioxide - quantitatively, approximately 65% as much warming as CO₂. And certain other short-lived species cause a cooling effect which offsets some of this warming.

This is not a new conclusion. The IPCC puts the [radiative forcing](#) from CO₂ at 1.66 W/m², compared to the forcing from other greenhouse gases, black carbon, and tropospheric ozone at approximately 1.4 W/m². Similarly, the negative forcing from aerosols is approximately -1.2 W/m².




Figure 1: Radiative forcing estimates from the [IPCC FAR](#)

Thus if anything, the 65% figure is an underestimate of the contributions of short-lived pollutants to global warming, but this contribution does not change the 1.66 W/m^2 radiative forcing from CO_2 or [the amount of global warming it has caused](#).

Much ado has also been made about another quote from the commentary:

"Warming over the past 100 years is consistent with high climate sensitivity to atmospheric carbon dioxide combined with a large cooling effect from short-lived aerosol pollutants, but it could equally be attributed to a low climate sensitivity coupled with a small effect from aerosols. These two possibilities lead to very different projections for future climate change."

This statement gets to the main point of the commentary - that there remains significant uncertainty regarding the effect of these short-lived pollutants on the global climate. However, estimates of the planetary climate sensitivity to increasing atmospheric CO_2 and other radiative forcings are not solely based on the change in the mean global temperature over the past 100 years. In fact, the climate sensitivity parameter has been estimated [through many different methods](#), including:

- climate models
- recent responses to large volcanic eruptions
- recent responses to solar cycles
- paleoclimate data
- data from the last Glacial Maximum
- and yes, data from the instrumental period

All of these different methods show strong agreement, overlapping in the IPCC climate sensitivity range of 2 to 4.5°C for a doubling of atmospheric CO_2 ($2\times\text{CO}_2$).




Figure 2: Distributions and ranges for climate sensitivity from different lines of evidence. The circle indicates the most likely value. The thin colored bars indicate very likely value (more than 90% probability). The thicker colored bars indicate likely values (more than 66% probability). Dashed lines indicate no robust constraint on an upper bound. The IPCC likely range (2 to 4.5°C) and most likely value (3°C) are indicated by the vertical grey bar and black line, respectively ([Knutti and Hegerl 2008](#))

Interestingly, Penner et al. find that whether the climate sensitivity parameter is on the low or high end, reducing anthropogenic emissions of the short-lived warming pollutants would achieve a significant reduction in global warming over the next 50-100 years. In the red lines in the Figure 3, they employ a climate model with a sensitivity of 5°C for 2xCO₂, slightly outside the IPCC likely range. The blue line is a climate model with a sensitivity of 2°C for 2xCO₂, on the lower end of the IPCC range. Note that even with the lower climate sensitivity, the model shows the planet warming 3°C by 2100 in this emissions scenario (see the figure caption for further details).




Figure 3: Global mean temperature measurements (black) and projections based on an IPCC scenario with high emissions (A2) for a climate sensitivity parameter of 5°C (upper red) and 2°C (upper blue). Linearly decreasing the total anthropogenic radiative forcing owing to methane, tropospheric ozone and black carbon — starting in 2010 and achieving pre-industrial levels by 2050 — results in significant near-term climate mitigation (lower blue and red curves) ([Penner 2010](#))

Unfortunately, reducing the short-lived cooling pollutants such as aerosols would cause a warming effect of similar magnitude, and so [CO₂ remains the primary pollutant of concern](#). Coincidentally, a group of scientists from NASA GISS just published a paper in *Science* entitled [Atmospheric CO₂: Principle Control Knob Governing Earth's Temperature](#).

Although it is important to reduce the remaining climate uncertainties, such as the magnitude of the impacts of short-lived pollutants, it does not change the fact that CO₂ is [very likely](#) the driving force behind the current global warming, or that if we double the amount of CO₂ in the atmosphere from pre-industrial levels, the planet will [likely](#) warm in the range of 2 to 4.5°C.



The Skeptical Science website by [Skeptical Science](#) is licensed under a [Creative Commons Attribution 3.0 Unported License](#).

 SkepticalScience.com



Skeptical Science explains the science of global warming and examines climate misinformation through the lens of peer-reviewed research. The website won the Australian Museum 2011 Eureka Prize for the Advancement of Climate Change Knowledge. Members of the Skeptical Science team have authored peer-reviewed papers, a [college textbook on climate change](#) and the book [Climate Change Denial: Heads in the Sand](#). Skeptical Science content has been used in university courses, textbooks, government reports on climate change, television documentaries and numerous books.



The [Skeptical Science](#) website by Skeptical Science is licensed under a Creative Commons [Attribution 3.0 Unported License](#).