



This is the print version of the [Skeptical Science](http://sks.to/albedo) article '[It's albedo](http://sks.to/albedo)', which can be found at <http://sks.to/albedo>.

The albedo effect and global warming

What The Science Says:

The long term trend from albedo is that of cooling. In recent years, satellite measurements of albedo show little to no trend.

Climate Myth: It's albedo

"Earth's Albedo has risen in the past few years, and by doing reconstructions of the past albedo, it appears that there was a significant reduction in Earth's albedo leading up to a lull in 1997. The most interesting thing here is that the albedo forcings, in watts/sq meter seem to be fairly large. Larger than that of all manmade greenhouse gases combined." ([Anthony Watts](#))

Change in the Earth's albedo is a potentially powerful driver of climate. When the planet's albedo or reflectivity increases, more sunlight is reflected back into space. This has a cooling effect on global temperatures. Conversely, a drop in albedo warms the planet. A change of just 1% to the Earth's albedo has a radiative effect of 3.4 Wm^{-2} , comparable to the forcing from a doubling of carbon dioxide. So how has albedo affected global temperatures in recent decades?

Albedo trends before 2000

There are various factors that affect the Earth's albedo. Snow and ice are highly reflective so when they melt, albedo drops. Forests have a lower albedo than open land so deforestation increases albedo (but for the record, no, chopping down all our forests is not the solution to global warming). Aerosols have a direct and indirect effect on albedo. The direct effect is reflecting sunlight back into space, cooling the Earth. The indirect effect is when aerosol particles act as a cloud condensation nucleus, affecting the formation and lifetime of clouds. Clouds in turn influence global temperatures in various ways. They cool the climate by reflecting incoming sunlight but can also warm the climate by trapping outgoing infrared radiation.

All these factors are considered when adding up the various radiative forcings that drive climate. Changes in land use are calculated from historical reconstructions of cropland and pastureland changes. Combinations of satellite and surface-based observations allow us to determine trends in aerosol levels as well as cloud albedo effect. What we observe is that of the various albedo forcings, cloud albedo is the most dominant effect. The long term trend is that of cooling with a radiative forcing from 1850 to 2000 of -0.7 Wm^{-2} .




Figure 1: Globally and annually averaged radiative forcing ([Chapter 2 of the IPCC AR4](#)).

Albedo trends after 2000

One way to measure the Earth's albedo is the use of earthshine. This is sunlight reflected from the Earth, then reflected from the Moon back to the nighttime Earth. Earthshine has been measured at the Big Bear Solar Observatory since November 1998 (with some measurements in 1994 and 1995). Figure 2 shows changes in albedo from reconstructed satellite data (black line) and Earthshine measurements (blue line) ([Palle 2004](#)).




Figure 2: Albedo anomalies reconstructed from ISCCP satellite data (black) and Earthshine-observed albedo anomalies (blue). The right hand vertical scale shows **negative** radiative forcing (eg - cooling) ([Palle 2004](#)).

The data in Figure 2 is problematic. The black line, reconstructed from ISCCP satellite data, "is a purely statistical parameter that has little physical meaning as it does not account for the non-linear relations between cloud and surface properties and planetary albedo and does not include aerosol related albedo changes such as associated with Mt. Pinatubo, or human

emissions of sulfates for instance" ([Real Climate](#)).

Even more problematic is the spike in albedo around 2003, shown by the blue earthshine line. This is in sharp contrast to satellite measurements which showed little to no trend over the same period. To put this in perspective, consider the Pinatubo volcanic eruption in 1991 which spewed aerosols into the atmosphere. These aerosols reflected incoming sunlight, causing a negative radiative forcing of 2.5 Wm^{-2} . This led to a dramatic drop in global temperatures. The earthshine data indicate a radiative forcing of nearly -6 Wm^{-2} which should cause an even greater drop in global temperatures. No such event occurred ([Wielicki 2007](#)).

In 2008, the reason for the discrepancy was discovered. The Big Bear Solar Observatory installed a new telescope in 2004 to measure earthshine. With the new and improved data, they recalibrated their old data and updated their earthshine albedo results ([Palle 2008](#)). Figure 3 shows the old albedo data (black) and the updated albedo (blue). The anomalous 2003 spike disappears. Nevertheless, a trend of increasing albedo remains from 1999 to 2003.




Figure 3: Earth albedo anomalies as measured by earthshine. In black are the albedo anomalies published in 2004 ([Palle 2004](#)). In blue are the updated albedo anomalies after improved data analysis, which also include more years of data ([Palle 2008](#)).

How accurate is the earthshine method in determining global albedo? The earthshine method doesn't give a global albedo estimate. It covers about one third of the Earth at each observation occasion and certain areas can never be "seen" from the measurement site. Furthermore the measurements are sparsely sampled in time, and only made in a narrow wavelength band of 0.4 to $0.7 \mu\text{m}$ ([Bender 2006](#)).

In contrast, satellite data such as CERES is a global measure of the Earth's reflected shortwave radiation, including the effects of all atmospheric and surface properties. It covers a broader spectrum than earthshine (0.3 - $5.0 \mu\text{m}$). An analysis of the CERES data finds no long term trend in albedo from March 2000 to June 2005. A comparison with 3 independent sets of satellite data (MODIS, MISR and SeaWiFS) also finds "remarkable consistency" between the 4 satellite results ([Loeb 2007a](#)).




Figure 4: Monthly anomalies in global mean CERES SW TOA flux and MODIS cloud fraction ([Loeb 2007b](#)).

Albedo has had an effect on global temperatures - mostly a cooling effect on long term trends. As for recent albedo trends, earthshine data shows increasing albedo from 1999 to 2003 but little to no trend from 2003. Satellites show little to no trend since 2000. The radiative forcing from albedo changes in recent years appears to be minimal.



Skeptical Science explains the science of global warming and examines climate misinformation through the lens of peer-reviewed research. The website won the Australian Museum 2011 Eureka Prize for the Advancement of Climate Change Knowledge. Members of the Skeptical Science team have authored peer-reviewed papers, a [college textbook on climate change](#) and the book [Climate Change Denial: Heads in the Sand](#). Skeptical Science content has been used in university courses, textbooks, government reports on climate change, television documentaries and numerous books.



The [Skeptical Science](#) website by Skeptical Science is licensed under a Creative Commons [Attribution 3.0 Unported License](#).