



This is the print version of the [Skeptical Science](http://sks.to/midcentury) article '[It cooled mid-century](http://sks.to/midcentury)', which can be found at <http://sks.to/midcentury>.

Why did climate cool in the mid-20th Century?

What The Science Says:

The average global surface temperature decreased slightly from 1940 to 1975. The main reason behind this mid-century cooling was global dimming due to anthropogenic sulfate aerosol emissions.

Climate Myth: It cooled mid-century

"It was the post war industrialization that caused the rapid rise in global CO2 emissions, but by 1945 when this began, the Earth was already in a cooling phase that continued until 1975. With 32 years of rapidly increasing global temperatures and only a minor increase in global CO2 emissions, followed by 33 years of slowly cooling global temperatures with rapid increases in global CO2 emissions, it was deceitful for the IPCC to make any claim that CO2 emissions were primarily responsible for observed 20th century global warming." ([Norm Kalmanovitch](#))

Mid-20th Century Cooling

There was a very slight cooling in the average global surface temperature from about 1940 to 1975. Although the global temperature only decreased by approximately 0.1°C, this period represents a divergence from the warming periods of 1915 to 1940 and 1975 to Present.




Figure 1: Global temperature anomaly and distinct linear trends over the 21st century ([Open Mind](#))

Climate scientists believe that the primary cause of this mid-century cooling was an increase in atmospheric aerosols due to anthropogenic emissions (primarily from the burning of fossil

fuels). Aerosols have a complex effect on the climate, because they have both direct and indirect impacts.

Direct Effect

The direct effect of aerosols on climate is the mechanism by which aerosols scatter and absorb shortwave and longwave radiation (a.k.a. "global dimming"), thereby altering the radiative balance of the Earth-atmosphere system. The key parameters for determining the direct aerosol radiative forcing are the aerosol optical properties and distribution in the atmosphere ([IPCC 2007](#)).

Indirect Effect

The indirect effect of aerosols on climate is the mechanism by which they modify the microphysical and, therefore, radiative properties, amount, and lifetime of clouds. A key parameter for determining the indirect effect of aerosols on the global surface temperature is the effectiveness of an aerosol particle to act as a cloud condensation nucleus - a function of the aerosol size, chemical composition, mixing state, and ambient environment ([IPCC 2007](#)).




Figure 2: Diagram of direct and indirect aerosols' radiative mechanisms ([IPCC 2007](#))

Radiative Forcing

We can attempt to quantify the impact of anthropogenic aerosol emissions on the average global temperature by first examining the associated radiative forcing. The radiative forcing is a measure of the influence that a factor has in altering the balance of incoming and outgoing energy in the Earth-atmosphere system.

According to the IPCC, the total (direct + indirect) radiative forcing due to anthropogenic aerosols could range anywhere from -0.4 to -2.7 Watts per square meter (W/m^2), although the most likely value is -1.2 W/m^2 .




Figure 3: Radiative forcing from various sources compared to 1750 (AR5 version)

The aerosol radiative forcing, particularly the indirect cloud albedo effect, is the largest source of uncertainty among all anthropogenic radiative forcings. However, keeping the range of possible values in mind, we can still estimate the cooling effect caused by these forcings during the mid-20th century. The majority of the increase in atmospheric sulfates occurred during this period:




Figure 4: Atmospheric sulfate aerosol concentration in parts per billion (ppb) from the GISP2 ice core (black) vs. anthropogenic sulfate emissions (red) since 1800 ([Open Mind](#))

The general natural dust baseline is around 30 ppb, with a rise to about 65 ppb in the early 20th century. Then from 1945 to 1970 (the mid-century period in question), atmospheric sulfate concentration increased from 65 to 120 ppb. This tells us that approximately 60% of the total

sulfate increase since 1800 occurred during the mid-century cooling period.

Direct Aerosol Forcing

The change in reflected solar flux is proportional to the optical thickness of the aerosol layer ([Charlson et al. 1991](#)).

$$\overline{\Delta F_R} \approx \frac{1}{2} Q_0 \overline{T_i^2} (1 - \overline{A_c}) (1 - \overline{R_s})^2 \beta \delta$$

where F_R is reflected solar flux, Q_0 is the solar constant (total solar irradiance), T_i is the atmospheric transmittance, A_c is the fractional coverage of clouds, β is the fraction of upward scattered flux, and δ is the optical thickness.

Using $Q_0 = 1366 \text{ W/m}^2$, $A_c = 0.61$ for globally averaged cloud cover, $T_i = 0.76$, and $R_s = 0.15$ (see [Charlson 1991](#) for references to the sources of these values), F_R is approximately 32 times the optical thickness δ . Charlson et al. estimate the optical thickness as the product of the aerosol burden and mass scattering coefficient, and find for cloudless skies, the optical thickness is approximately 2.8×10^{-2} . Thus from the formula above, the reflected solar flux is approximately 0.9 W/m^2 , which is on the high end of the IPCC range of values for the direct aerosol forcing.

The indirect sulfate effect on cloud formation is more complex, but we'll proceed under the assumption that the total sulfate radiative forcing is approximately proportional to the atmospheric sulfate aerosol concentration, in order to come up with a ballpark estimate regarding its impact on mid-century cooling. In other words, we'll assume that a 60% increase in atmospheric sulfate concentration corresponds to a 60% increase in radiative forcing. Also note that sulfates are not the only anthropogenic aerosols, but do comprise the majority of the radiative forcing.

Approximation of sulfate aerosol cooling effect

With these assumptions in mind, approximately 60% of the -0.4 to -2.7 W/m^2 net aerosol forcing occurred during mid-century, or -0.24 to -1.62 W/m^2 , with a most likely value of -0.72 W/m^2 . To convert this to a surface temperature change, we need to multiply by the [climate sensitivity](#) factor (λ), which is 0.54 to $1.2^\circ\text{C}/(\text{W/m}^2)$ ([IPCC 2007](#)). The relationship between surface temperature change (dT) and radiative forcing (dF) is:

$$dT = \lambda * dF = (0.54 \text{ to } 1.2) * (-0.24 \text{ to } -1.62) = -0.13 \text{ to } -1.9^\circ\text{C} \text{ with a most likely value of } -0.58^\circ\text{C}$$

We can also compare this to the [temperature change caused by anthropogenic CO2](#) during the period in question:

$$dT = \lambda * dF = (0.54 \text{ to } 1.2) * 5.35 * \ln(331/308) = 0.21 \text{ to } 0.46^\circ\text{C} \text{ with a most likely value of } 0.31^\circ\text{C}$$

In addition to CO_2 , other anthropogenic greenhouse gas emissions and solar output during this period caused an additional $\sim 0.15^\circ\text{C}$ of warming. Therefore, using the most likely values calculated above, we would expect to see an approximately 0.1°C cooling of global average surface temperatures from 1940 to 1975. And indeed that is what we observe:




Figure 5: Average global land and ocean surface temperature ([NASA GISS](#))

Days cooled, but nights warmed

Another signature of aerosol cooling offsetting greenhouse gas warming is the continued increase in daily minimum, nighttime temperatures while the maximum, daytime temperatures drop. This is because aerosols cause global dimming by scattering sunlight; an effect which is much more influential during the day when solar radiation is bombarding the Earth's surface. At night, greenhouse gases continue to absorb and re-radiate thermal radiation from the Earth's surface, which causes the nighttime surface temperatures to continue warming despite the cooling daytime temperatures due to the aerosol dimming.

[Wild et al. \(2007\)](#) investigated these effects and found that from 1958 to 1985, daytime land surface temperatures cooled at a rate of -0.04°C per decade, but nighttime temperatures warmed at a rate of 0.11°C per decade. From 1982 to 2002, they found that daytime land surface temperatures warmed at a rate of 0.37°C per decade, while nighttime temperatures warmed by 0.40°C per decade.

The fact that nighttime temperatures continued to warm while daytime temperatures cooled mid-century is a strong indicator of the combined effects of anthropogenic aerosol and greenhouse gas emissions, while the increased rate of warming over the past 3 decades reflects the increasing atmospheric concentration and radiative forcing from greenhouse gases.

What happened since 1975?

Clearly since about 1975, global surface temperatures have trended rapidly upwards (at a rate of nearly 0.2°C per decade). So what caused the mid-century cooling to end?

The main cause of the sudden shift in global temperature trends was the passage of [Clean Air Acts](#) by various countries in response to air pollution and acid rain. The USA, for example, first passed its [Clean Air Act in 1970](#), with amendments in 1977 and 1990. Coincidentally, the US Supreme Court (in [Massachusetts v. EPA](#)) and EPA (in an [endangerment finding](#)) also recently decided that greenhouse gases qualify as 'air pollutants' in the Clean Air Act and must be regulated accordingly.

Under the Clean Air Acts, sulfate emissions were regulated, and as a consequence their rapid atmospheric increase was stabilized right around 1975:




Figure 6: Global anthropogenic sulfur emissions ([Pacific Northwest National Labs](#))

Meanwhile anthropogenic greenhouse gas emissions have continued to increase. Since 1975, the atmospheric CO₂ concentration alone has increased from 331 to 392 parts per million by volume, which corresponds to a temperature increase of about 0.7°C, though we've only seen about 0.55°C warming over that period due to the thermal inertia of the oceans and ['warming in the pipeline'](#).

Mid-century cooling was primarily anthropogenic

To sum up, anthropogenic sulfur emissions appear to be the main cause of the mid-century cooling. These emissions decreased the mean global surface temperature by approximately 0.5°C during this period, while anthropogenic greenhouse gas emissions caused a warming of approximately 0.4°C. Therefore, even though greenhouse gas emissions continued to have a warming effect during this period, it was more than offset (hidden) by anthropogenic aerosol emissions, until those emissions were brought under control by government intervention while greenhouse gas emissions continued to increase unabated. In other words, the mid-century cooling is actually an expected result based on our current understanding of climate science, and is successfully hindcasted by climate models ([Meehl 2004](#)).




Figure 7: Anthropogenic plus natural vs. just natural radiative forcing temperature change vs.

observed global surface temperature increase ([Meehl 2004](#))

However, the overall impact of sulfate aerosols, particularly due to their indirect effects via cloud formation, remain a significant source of uncertainty. Despite this uncertainty, they remain the likely dominant cause of the slight mid-20th century cooling.



Skeptical Science explains the science of global warming and examines climate misinformation through the lens of peer-reviewed research. The website won the Australian Museum 2011 Eureka Prize for the Advancement of Climate Change Knowledge. Members of the Skeptical Science team have authored peer-reviewed papers, a [college textbook on climate change](#) and the book [Climate Change Denial: Heads in the Sand](#). Skeptical Science content has been used in university courses, textbooks, government reports on climate change, television documentaries and numerous books.



The [Skeptical Science](#) website by Skeptical Science is licensed under a Creative Commons [Attribution 3.0 Unported License](#).